



Nutritional genomics and metabolomics in obesity and type 2 diabetes

Citation

Hu, Frank B, and Kothandaraman Narasimhan. 2014. "Nutritional genomics and metabolomics in obesity and type 2 diabetes." BMC Genomics 15 (Suppl 2): 010. doi:10.1186/1471-2164-15-S2-010. <http://dx.doi.org/10.1186/1471-2164-15-S2-010>.

Published Version

doi:10.1186/1471-2164-15-S2-010

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:12717407>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

ORAL PRESENTATION

Open Access

Nutritional genomics and metabolomics in obesity and type 2 diabetes

Frank B. Hu^{1,2*}, Kothandaraman Narasimhan²

From 2nd International Genomic Medical Conference (IGMC 2013)
Jeddah, Kingdom of Saudi Arabia. 24-27 November 2013

Obesity and type-2 diabetes arise from the complex interplay of both genetic and environmental factors. In the past several years, we have seen considerable progress in the use of genomics and metabolomics in investigating the role of diet and nutrition in the etiology of obesity and diabetes. Recent work suggests that the adverse effect of several obesity loci especially FTO may be modified by dietary factors. In a recent study (Qi et al. NEJM 2012), we found a significant interaction between consumption of Sugar-Sweetened Beverages (SSBs) and genetic variants related to Body Mass Index (BMI) in three cohort studies. A genetic-predisposition score was calculated on the basis of 32 BMI-associated loci. We found that the genetic association with BMI was more pronounced among participants with higher intake of SSBs than among those with lower intake. In the combined cohorts, the increases in BMI per increment of 10 risk alleles were 1.00 unit for an intake of < 1 serving/month, 1.12 for 1-4 servings/month, 1.38 for 2-6 servings/week, and 1.78 for ≥ 1 servings/day ($P < 0.001$ for interaction). This analysis suggests that the genetic association with adiposity is amplified by greater intakes of SSBs. In the area of nutrition and metabolomics, several recent epidemiological studies have found that plasma concentrations of metabolites especially Branched-Chain Amino Acids (BCAAs) are associated with increased risk of type-2 diabetes. Other novel metabolite classes have also been linked to diabetes, such as short- and medium-chain acylcarnitines, the specific lipid classes of sphingomyelins (SMs), lysophosphatidylcholines, phosphatidylcholines (PCs), and lysophosphatidylethanolamines. Metabolomics has also been used to characterize the complex human metabolic effects of specific foods, nutrients, and dietary patterns in both intervention and epidemiological studies. Our

preliminary analysis found that urinary levels of gut flora metabolites derived from dietary intakes of polyphenols were significantly associated with risk of diabetes.

In summary, both nutrigenomics and metabolomics studies have provided new insights to the etiology of obesity and diabetes and individual differences in response to diet. Continued technological advances in sensitive high-throughput methods and enhanced bioinformatics and analytical tools in combination with carefully conducted population-based studies will enable more widespread use of these technologies in nutrition and metabolic disease research, which will eventually help to achieve the goal of personalized nutrition for prevention and treatment of chronic diseases.

Authors' details

¹Departments of Nutrition and Epidemiology, Harvard School of Public Health, Boston, MA, USA. ²Center of Excellence in Genomic Medicine Research, King Abdulaziz University, Jeddah, KSA.

Published: 2 April 2014

doi:10.1186/1471-2164-15-S2-O10

Cite this article as: Hu and Narasimhan: Nutritional genomics and metabolomics in obesity and type 2 diabetes. *BMC Genomics* 2014 15(Suppl 2):O10.

¹Departments of Nutrition and Epidemiology, Harvard School of Public Health, Boston, MA, USA

Full list of author information is available at the end of the article